

Supplementary Table 17. Marginal predicted values of the fixed and random-effects model for the changes from baseline in pulp/tooth proportion values ($\Delta B1/B0$) in the DPC, PP, and CP groups against those in the LB group

Treatment	Follow-up period										
	6 mo	12 mo	18 mo	24 mo	30 mo	36 mo	48 mo	60 mo	72 mo		
LB	0.52 (0.34–0.71)	0.94 (0.67–1.21)	1.32 (0.96–1.68)	1.66 (1.21–2.11)	1.96 (1.42–2.51)	2.23 (1.59–2.86)	2.63 (1.81–3.45)	2.88 (1.88–3.89)	2.97 (1.78–4.17)		
DPC	0.59 (0.42–0.76)	1.02 (0.76–1.27)	1.38 (1.03–1.72)	1.68 (1.24–2.11)	1.91 (1.39–2.43)	2.08 (1.47–2.69)	2.23 (1.44–3.02)	2.12 (1.16–3.09)	1.76 (0.62–2.91)		
PP	0.55 (0.42–0.68)	1.01 (0.82–1.20)	1.42 (1.17–1.68)	1.79 (1.47–2.11)	2.11 (1.72–2.49)	2.38 (1.93–2.82)	2.77 (2.20–3.35)	2.98 (2.27–3.69)	3.00 (2.16–3.85)		
CP	0.72 (0.55–0.89)	1.37 (1.12–1.62)	1.97 (1.64–2.31)	2.53 (2.11–2.95)	3.05 (2.55–3.56)	3.53 (2.93–4.14)	4.36 (3.53–5.20)	5.02 (3.89–6.15)	5.51 (3.99–7.03)		

Numbers in parentheses are 95% confidence intervals.

CP, coronal pulpotomy; DPC, direct pulp capping; LB, protective liner and base; PP, pulpotomy.