

Supplementary Table 32. Marginal predicted values of the fixed and random-effects model for the changes from baseline of pulp/tooth proportion values ($\Delta B2/B0$) in the different groups of preoperative pulpal diagnosis, against the normal pulp group

Preoperative diagnosis	Follow-up period									
	6 mo	12 mo	18 mo	24 mo	30 mo	36 mo	48 mo	60 mo	72 mo	
Normal pulp	0.62 (0.19 to 1.04)	1.20 (0.54 to 1.86)	1.70 (0.80 to 2.61)	2.12 (0.97 to 3.27)	2.46 (1.06 to 3.86)	2.72 (1.07 to 4.37)	3.00 (0.85 to 5.15)	2.96 (0.31 to 5.60)	2.59 (-0.56 to 5.74)	
Reversible pulp	0.75 (0.46 to 1.03)	1.36 (0.91 to 1.81)	1.91 (1.29 to 2.52)	2.38 (1.59 to 3.16)	2.78 (1.82 to 3.74)	3.11 (1.99 to 4.24)	3.56 (2.10 to 5.03)	3.73 (1.92 to 5.54)	3.62 (1.47 to 5.77)	
Irreversible pulp	1.31 (1.08 to 1.55)	2.27 (1.91 to 2.63)	3.15 (2.66 to 3.65)	3.97 (3.35 to 4.60)	4.73 (3.96 to 5.49)	5.41 (4.51 to 6.31)	6.59 (5.42 to 7.76)	7.49 (6.05 to 8.94)	8.13 (6.41 to 9.86)	

Numbers in parentheses are 95% confidence intervals.